


Photoelectric absorption as an x-ray spectral diagnostic of massive star winds


Massive star winds



The winds of massive stars deposit chemically-enriched matter as well as energy and momentum into their galactic environment (wind-blown bubble NGC 7635 at left). The x-rays produced in the spatially-unresolved circumstellar environment are a sensitive diagnostic of the mass-loss rate, the key measurable parameter that determines the influence of stellar winds. Typical mass-loss rates of O stars are $\dot{M} \sim 10^{-6} M_{\odot}/\text{yr}$.


X-ray production

Instabilities lead to shock-heating of the wind. These shocked regions, which make up a small fraction of the total wind mass, reach temperatures between 10^6 and 10^7 K. In the simulation shown below, shocks start developing about half a stellar radius above the surface of the star.



Theoretical predictions


1. In the shock regions, highly ionized atoms produce thermal x-ray emission lines.
2. Production in the fast-moving wind results in Doppler-broadened lines.
3. **The cool material in the wind absorbs x-rays.** We should find evidence of this absorption in both the overall spectrum and in the shape of individual line profiles.




Erin Martell,¹ Emma Wollman,¹ David Cohen,¹ Marc Gagné,² and Maurice Leutenegger³

1: Swarthmore College; 2: West Chester University
3: NASA/Goddard Space Flight Center

Chandra data



Wind absorption models



Chandra spectra of massive stars

High-resolution x-ray spectra of massive stars show a trend in spectral hardness with stellar surface temperature (and mass and luminosity): above left, the hottest stars at the top have the hardest spectra.

Is this trend due to the emission temperature of shock-heated plasma?

Or, is it due to increasing **wind absorption** of x-rays for hotter stars that have stronger winds?


We have developed a radiation transport model of thermal x-ray emission from plasma embedded within an extended stellar wind. Preliminary models that all assume the same intrinsic spectrum but with differing amounts of wind attenuation (shown in the right panel) reproduce the observed hardness trend. There may still be some trend in the temperature distribution, however – ratios of H/He-like lines are the key to determining this (see the highlighted Mg XII/XI lines in the left panel).

Atomic opacity

The opacity of the cold, unshocked component of the wind is due to K-shell photoionization of low-Z elements: C, N, O, Ne, Mg and the L-shell of Fe and Ni in a cosmic abundance mixture.


The opacity is greater at longer wavelengths: Wind absorption hardens the overall x-ray spectrum (figure at top of this column).

And this continuum opacity also skews the individual Doppler-broadened line profiles (explained in the right-hand column).




Wind geometry

The shape of the individual x-ray emission lines is also governed by absorption. The diagram to the right shows contours of constant radial velocity as seen by an observer to the left. Red-shifted x-rays emitted from the back side of the wind must pass through more material than blue photons emitted from the front of the wind. Red photons are thus preferentially absorbed, leading to asymmetric lines.




Line asymmetry

In our model of the line profile shape, the asymmetry is parameterized by τ_* , the fiducial optical depth. τ_* is defined as $\kappa \dot{M} / 4\pi R_* v_{\infty}$, where κ is the opacity, \dot{M} is the mass-loss rate, R_* is the stellar radius, and v_{∞} is the terminal velocity. Plotted to the right are four un-normalized models with different values of τ_* .



Fitting the data



The Fe XVII line at 15.014 angstroms in ζ Pup is shown with two models. The best-fit curve (dark blue) is clearly less shifted than predicted by the H α mass-loss rate (light blue curve), indicating a lower mass-loss rate. The fit also gives us the radius of x-ray emission onset, which is 0.55 stellar radii above the star's surface for this line. This result is consistent with

the hydrodynamical simulation shown in the left column of this poster.

Mass-loss rate determination from global fit

The Fe XVII line is not the only line in ζ Pup's spectrum to show less asymmetry than the H α mass-loss rate would require. Using the wavelength-dependent opacity, we can fit a single mass-loss rate to all lines in the spectrum. We see that the H α mass-loss rate (light blue) is inconsistent with the data and that a lower mass-loss rate (dark blue) is preferred. In other stars, the best-fit mass-loss rates are up to an order of magnitude lower than values obtained from other diagnostics. **Such a reduction in mass-loss rates would affect our understanding of stellar and galactic evolution.**

